

# Abundances in Damped Ly $\alpha$ Galaxies

Paolo Molaro<sup>1</sup>

Osservatorio Astronomico di Trieste-INAf Via G.B. Tiepolo 11,  
I-33100 Trieste, Italy

**Abstract.** Damped Ly $\alpha$  galaxies provide a sample of young galaxies where chemical abundances can be derived throughout the whole universe with an accuracy comparable to that for the local universe. Despite a large spread in redshift, HI column density and metallicity, DLA galaxies show a remarkable uniformity in the elemental ratios rather suggestive of similar chemical evolution if not of an unique population. These galaxies are characterized by a moderate, if any, enhancement of  $\alpha$ -elements over Fe-peak elemental abundance with  $[S/Zn] \approx 0$  and  $[O/Zn] \approx 0.2$ , rather similarly to the dwarfs galaxies in the Local Group. Nitrogen shows a peculiar behaviour with a bimodal distribution and possibly two plateaux. In particular, the plateau at low N abundances ( $[N/H] < -3$ ), is not observed in other astrophysical sites and might be evidence for primary N production by massive stars.


## 1 Introduction

Any slab of intervening material along the line of sight of a background source with hydrogen column density high enough to produce damping wings in Ly $\alpha$ , conventionally  $\log N(HI) > 20.3 \text{ cm}^{-2}$ , is producing a Damped Ly $\alpha$  galaxy (DLA). Damped systems hold a large fraction of neutral gas at high redshift and are considered the progenitors of present day galaxies. Holding neutral gas and being free from ionization effects, DLAs provide column densities at very high precision ( $\approx 10\%$ ), which together with the fact that they are observed up to redshift  $\approx 4.5$  or equivalently at a look-back time of  $\approx 12$  Gyr, expand to almost the entire universe the possibility of a detailed chemical investigation. Observed metallicities are generally low, with  $[Fe/H]$  varying between  $-2.5$  and  $-1.0$ , but never below  $-2.5$  and with a mild evolution with redshift. The elemental abundances, which we discuss here in more detail, resemble very closely those observed in the dwarf and irregular galaxies of the Local Group which are the subject of this conference.

## 2 Dust

Dust is probably present in the DLA and significantly affects the observed abundances. I counted 55 systems for which both Fe and Zn are measured, which are plotted in Fig. 1. The abundance of Fe is always found below that of Zn. By analogy with the interstellar medium, this behaviour is interpreted as the effect of some Fe being locked into dust grains. Other indicators for the presence of dust

are the correlation of  $[\text{Fe}/\text{Zn}]$  with  $\text{H}_2$  and the reddening excess of QSOs behind Damped galaxies. The figure shows a clear trend of Fe depletion with metallicity. There is also evidence for a sort of threshold line  $[\text{Fe}/\text{Zn}] \approx -(2 + [\text{Zn}/\text{H}])$  with no Fe depletion below it, which points out to the presence of a forbidden region for dust formation. This behaviour may be relevant for the understanding of dust formation in low metallicity environments.




**Fig. 1.** Dust evidence in the DLAs


### 3 The $[\alpha/\text{Zn}]$ ratio

The presence of dust affects abundances of elements participating into dust grains, and it needs either to be corrected or to be avoided. Considering the uncertainties involved in dust formation, the second approach looks safer. S and O are  $\alpha$  elements showing little affinity with dust, while Zn is, rather uncomfortably, the only one available among the iron-peak group. In Fig 2 the  $[\text{S}/\text{Zn}]$  elements for 24 Damped galaxies are shown. The average value is consistent with zero:  $\langle [\text{S}/\text{Zn}] \rangle = 0.01 \pm 0.14$ . There is also a hint of decreasing  $[\text{S}/\text{Zn}]$  with the increasing of metallicity. O has been measured in a limited number of objects and those in which also Zn is available are plotted in Fig 3, together with a couple of particularly low  $[\text{O}/\text{Fe}]$ , which are a significant upper bound to the  $[\text{O}/\text{Zn}]$  value. QSO 0347-3819 is generally considered the best case for evidence of  $\alpha$  enhancement. However, the figure shows a new  $[\text{O}/\text{Zn}]$  measure from a revised measurement of ZnII by Levshakov et al ( $\log N(\text{ZnII}) = 12.26 \text{ cm}^{-2}$ , 2004, private com.). The revised ratio is  $[\text{O}/\text{Zn}] = 0.1 \pm 0.1$  and there is no more strong  $\alpha$  enhancement in this DLA. The case of Q 0812+32, with  $[\text{O}/\text{Zn}] = 0.35$ , should be probably considered separately since this DLA is rather peculiar and is considered the progenitor of an elliptical. Thus, with the exception of the QSO

0812+32 case, all the DLA galaxies show mild or absent  $\alpha$  enhancement, reminiscent of what found in the stars of the Local Group galaxies. To increase the statistical significance of the result one can use Si that, despite being refractory, tracks S rather well on average. In 40 DLA galaxies  $[\text{Si}/\text{Zn}] \approx 0$ , with only few exceptions of particularly dusty objects, showing that the absence of  $\alpha$  enhancement is not due to poor statistics but is a rather general property as was first suggested by Molaro et al (1996)



**Fig. 2.**  $[\text{S}/\text{Zn}]$  in DLAs




**Fig. 3.**  $[\text{O}/\text{Zn}]$  in DLAs, with some  $[\text{O}/\text{Fe}]$

### 3.1 Nitrogen

N shows little affinity with dust and is thought to be produced by intermediate mass stars in the 4-8  $M_{\odot}$  range which undergo HBB in the AGB phase. N has a dual nature since it is primary at low metallicities and secondary at high metallicities, as observed in the extragalactic HII regions. Being produced by relatively small masses it is produced with a time delay compared to O or other  $\alpha$  a fact that can explain the rather scattered  $[\text{N}/\text{Si}]$  at low abundances. It has been suggested by Prochaska et al (2002) that the DLAs show a bimodal distribution rather than a pure scatter behaviour. From the 2 systems initially considered by them there are now 5 systems and 2 upper limits that support the proposal for a bimodal distribution. The existence of two different regimes can be better appreciated when  $[\text{N}/\text{Si}]$  is plotted versus  $[\text{N}/\text{H}]$  as shown in Fig. 4. The plot clearly shows that the in DLA low and high  $[\text{N}/\text{Si}]$  occupy two separate regions. A possible explanation is that the low values are young systems where N comes from massive stars and the AGB have not yet started to produce the bulk of N. A primary N production by massive stars is not foreseen by standard models but it is present in the zero metallicity models of Chieffi & Limongi (2002) and the predicted  $[\text{N}/\text{Si}]$  ratios after integration over an appropriate IMF show ratios close to the observed ones. It would be rather appealing to find this N as the *smoking gun* of zero metal stars. In Fig 4 the  $[\text{N}/\text{Si}]$  of DLAs are shown together with the recent N determinations provided in halo stars. Stellar values

are in rough agreement with this picture, showing no values below  $[N/Si]=-1.5$ , although they populate also the portion of the diagram with  $[N/H] < -3.0$ .



**Fig. 4.** Nitrogen in DLAs, circles, and halo stars from Spite et al (2004), losange, and from Israelian et al (2004), squares.

### 3.2 A single population?

Determination of relative abundances of elements produced differently by Type I, TypeII SNa and IMS probes the early chemical build-up. Chemical evolutionary models characterized by a relatively low star formation rate have been proposed to explain the mild  $[\alpha/Zn]$  and the  $[N/Si]$  ratios in the DLA. A low SFR is commonly found in dwarfs, irregulars, LSB galaxies or in the external part of rotational disks. Observationally the DLAs show a remarkable constancy in the observed elemental ratios. So far there are 15 DLAs for which we have the whole triplet of elements N, Si and Zn. Si in this case is used as a fair proxy for O or S. The observed elemental ratios are:  $\langle [S/Zn] \rangle = 0.01 \pm 0.18$  and  $\langle [N/Si] \rangle = -0.79 \pm 0.16$ , i.e. remarkably uniform considering that the metallicity in this sample is varying from -2.0 to -0.5, redshift from 1.6 to 3.4 and  $N(HI)$  from 20.0 to 21.5. Such a homogeneity in the ratios is strongly suggestive of very similar evolutionary properties of the DLAs if not of a specific population of galaxies, which might dominate the DLA class at least at high redshift.

## References

1. A. Chieffi, M. Limongi Ap.J. **577**, 281 (2002)
2. G. Israelian, A Ecuivillon et al: A&A **421**, 649 (2004)
3. P. Molaro, S. D’Odorico et al: A&A **308**, 1 (1996)
4. J. Prochaska, R.B.C. Henry et al PASP **114**, 1197 (2002)
5. M. Spite, R. Cayrel et al: astroph-0409536 (2004)